

Reply to a comment by Peres et al.

G. Baskaran¹ and S.A. Jafari^{1,2}

¹Institute of Mathematical Sciences, C.I.T. Campus, Chennai 600 113, India

²Department of Physics, Sharif University of Technology, Tehran 11365-9161, Iran

Peres et al. comment [1] that in our recent work on graphite [2] we get spin-1 collective (spin-1 zero sound) mode only when we use, in addition to ‘random phase approximation’ (RPA), another approximation.

To justify our method and also clarify the issue involved, **particularly at low energies**, we discuss our spin-1 zero sound in a fermi liquid theory framework [3] and emphasize an important aspect of Landau’s zero sound collective mode analysis. One calculates the two particle forward scattering amplitude defined by Landau $\Gamma(p_1, p_2; k)$ by focusing on the singularity in the zero sound channel. Most importantly, as emphasized by Landau [4] one considers intermediate states in which the number of ‘quasi-particles’ are not changed. The RPA susceptibility we have computed in our approximation precisely corresponds to a bubble sum of such a process, where the quasi-particle number is conserved in the intermediate states.

To elaborate this point we write the ‘bare’ Hubbard Hamiltonian for the ideal graphene sheet in terms of the conduction band electron (e ’s) and valence band hole (h ’s) operators with **positive energies**:

$$\begin{aligned}
H = & \sum_{k\sigma} \epsilon_k (e_{k\sigma}^\dagger e_{k\sigma} + h_{k\sigma}^\dagger h_{k\sigma}) \\
& + \frac{U}{N} \sum_{kk'} \gamma_{k,k',q}^{(1)} (e_{k\uparrow}^\dagger e_{k+q\uparrow} h_{k'\downarrow}^\dagger h_{k'-q\downarrow} + \dots) \\
& + \frac{U}{N} \sum_{kk'} (\gamma_{k,k',q}^{(2)} (e_{-k\uparrow}^\dagger h_{k+q\uparrow}^\dagger e_{-k'\downarrow}^\dagger h_{k'-q\downarrow}^\dagger \\
& + \gamma_{k,k',q}^{(3)} e_{k\uparrow}^\dagger e_{k-q\uparrow}^\dagger e_{-k'\downarrow}^\dagger h_{k'-q\downarrow}^\dagger + \dots)
\end{aligned} \tag{1}$$

Here $\epsilon_k \equiv t\sqrt{1 + 4\cos\frac{\sqrt{3}k_x}{2}\cos\frac{k_y}{2} + 4\cos^2\frac{k_y}{2}}$ and γ ’s are complex form factors, $|\gamma| \sim 1$. Hubbard interaction terms describe various two body processes: electron-electron, hole-hole, electron-hole scattering, electron decay, vacuum polarization etc. The second line of eqn.(1), $e^\dagger e h^\dagger h$ is the one important for us - it represents scattering between a real electron and a real hole. The term $e^\dagger e^\dagger h^\dagger h^\dagger$ of the third line corresponds to spontaneous creation of two electron-hole pairs out of the vacuum and the second term $e^\dagger e h^\dagger e^\dagger$ represents decay of a real electron into an electron and electron-hole pair. That is, all terms contained in the third line are either processes that renormalize the vacuum or dress the quasi-particles. At the end, if a fermi liquid state is formed, one has ‘stable’ low energy renormalized **quasi particles that neither feel the vacuum polarization processes nor polar-**

ize the vacuum. In a renormalization group sense, the $\gamma^{(2)}, \gamma^{(3)}$ etc. terms of eqn.(1) are expected to renormalize to zero, as one approaches a fermi liquid fixed point. For our analysis of spin-1 zero sound in graphite, assumed to be a fermi liquid, we are essentially left with the $\gamma^{(1)}$ term with a renormalized $\tilde{U} > 0$. That is, one low energy electron and one hole quasi-particle repeatedly scatter, in a renormalized **frozen fermi liquid vacuum**, without intervention of any additional quasi-particles in the intermediate state, in the way described by Landau. This leads to our simple susceptibility expression [2]

$$\chi(q, \omega) = \frac{\chi_0(q, \omega)}{1 - \tilde{U}\chi_0(q, \omega)} \tag{2}$$

that has poles corresponding to our predicted spin-1 collective mode branch, below the particle-hole continuum.




FIG. 1. A diagram that is contained in Peres et al.’s bubble summation. Intermediate state A is a 1-electron and 1-hole state; B is a 3-electron and 3-hole state. At low energies, in a Landau theory such diagrams with more than 1-electron and 1-hole in the intermediate states do not exist, because $U\gamma^{(2)}$ is renormalized to zero/low values.

Peres et al., have, on the other hand, calculated a ‘2-sub lattice susceptibility tensor’, which uses the full unrenormalized Hubbard interaction terms [lines 2,3 and 4 in eqn.(1)] in their bubble summation procedure. When converted into a physically more transparent valence and conduction band quasi-particle basis of our fermi liquid state, Peres et al.’s susceptibility contains diagrams in which intermediate states have higher quasi-particle numbers (an example is shown in figure 1). Once we **isolate and remove these processes from their bubble summation**, we recover our fermi liquid result (equation 2) and a full spin-1 collective mode branch emerges from the bottom of the particle-hole continuum.

[1] N.M.R. Peres, M.A.N. Araujo and A.H. Castro Neto, preceding comment

- [2] G. Baskaran and S.A. Jafari, Phys. Rev. Lett. **89** 016402 (2002)
- [3] G. Baskaran and S.A. Jafari, unpublished
- [4] L.D. Landau, Sov. Phys. JETP, **35** 70 (1959); reprinted in 'The Many-Body Problem' Ed., D. Pines (W.A. Benjamin inc.) paragraph following eqn.(27) in page 277.